

		Vigamox (Moxifloxacin)	PredForte (Prednisolone)	Nevanac (Nepafenac)
Day Before Surgery	AM	1 Drop ____	1 Drop ____	1 Drop ____
	MIDDAY	1 Drop ____	1 Drop ____	1 Drop ____
	AFTERNOON	1 Drop ____	1 Drop ____	1 Drop ____
	EVENING	1 Drop ____	1 Drop ____	1 Drop ____
Day Of Surgery	AM	1 Drop ____	1 Drop ____	1 Drop ____
	SURGERY	SURGERY	SURGERY	SURGERY
		1 Drop every hour until bedtime	1 Drop - 4 times spread across the rest of the day	1 Drop - 4 times spread across the rest of the day
Day 1	AM	1 Drop ____	1 Drop ____	1 Drop ____
	MIDDAY	1 Drop ____	1 Drop ____	1 Drop ____
	AFTERNOON	1 Drop ____	1 Drop ____	1 Drop ____
	EVENING	1 Drop ____	1 Drop ____	1 Drop ____
Day 2	AM	1 Drop ____	1 Drop ____	1 Drop ____
	MIDDAY	1 Drop ____	1 Drop ____	1 Drop ____
	AFTERNOON	1 Drop ____	1 Drop ____	1 Drop ____
	EVENING	1 Drop ____	1 Drop ____	1 Drop ____
Day 3	AM	1 Drop ____	1 Drop ____	1 Drop ____
	MIDDAY	1 Drop ____	1 Drop ____	1 Drop ____
	AFTERNOON	1 Drop ____	1 Drop ____	1 Drop ____
	EVENING	1 Drop ____	1 Drop ____	1 Drop ____
Day 4	AM	1 Drop ____	1 Drop ____	1 Drop ____
	MIDDAY	1 Drop ____	1 Drop ____	1 Drop ____
	AFTERNOON	1 Drop ____	1 Drop ____	1 Drop ____
	EVENING	1 Drop ____	1 Drop ____	1 Drop ____
Day 5	AM	1 Drop ____	1 Drop ____	1 Drop ____
	MIDDAY	1 Drop ____	1 Drop ____	1 Drop ____
	AFTERNOON	1 Drop ____	1 Drop ____	1 Drop ____
	EVENING	1 Drop ____	1 Drop ____	1 Drop ____
Day 6	AM	1 Drop ____	1 Drop ____	1 Drop ____
	MIDDAY	1 Drop ____	1 Drop ____	1 Drop ____
	AFTERNOON	1 Drop ____	1 Drop ____	1 Drop ____
	EVENING	1 Drop ____	1 Drop ____	1 Drop ____
Day 7	AM	1 Drop ____	1 Drop ____	1 Drop ____
	MIDDAY	1 Drop ____	1 Drop ____	1 Drop ____
	AFTERNOON	1 Drop ____	1 Drop ____	1 Drop ____
	EVENING	1 Drop ____	1 Drop ____	1 Drop ____
		Discontinue after 1 week	Continue same drop schedule until bottle empty	Continue same drop schedule until bottle empty

****Ensure you wait five minutes after administering an eye drop before putting in another one to allow for proper absorption.****